

## IDENTIFICATION AND CHARACTERIZATION OF AN ANTHOCYANIN MUTANT (*ANT1*) IN TOMATO

### FIELD OF THE INVENTION

The present invention relates to a plant phenotype, designated *Anthocyanin 1* (*ANT1*), together with DNA and polypeptide sequences associated with the same.

### BACKGROUND OF THE INVENTION

Flavonoids comprise a diverse collection of red to blue colored secondary metabolites that accumulate in the tissues of many plant species. The primary structure of flavonoids consists of two aromatic carbon groups; benzopyran (A and C rings) and benzene (B ring). The variation in the heterocyclic C-ring of flavonoids and the interlinkage between the benzopyran and benzene groups are the basis for the classification of flavonoids into the flavone, flavonol, flavonone, isoflavone, anthocyanin, and flavane groups.

Anthocyanins have been associated with many important physiological and developmental functions in the plants, including, modification of the quantity and quality of captured light (Barker DH et al., *Plant Cell and Environment* 20: 617-624, 1977.); protection from the effects of UV-B radiation (Burger J and Edwards GE. *Plant and Cell Physiology* 37: 395-399, 1996; Klaper R et al., *Photochemistry and Photobiology* 63: 811-813, 1996); defense against herbivores (Coley and Kusar. In: Mulkey SS, Chazdon RL, Smith AP, eds. *Tropical Forest Plant Ecophysiology*. New York: Chapman and Hall 305-335, 1996); and protection from photoinhibition (Gould KS, et al., *Nature* 378: 241-242, 1995; and Dodd IC et al., *Journal of Experimental Botany* 49: 1437-1445, 1998); and scavenging of reactive oxygen intermediates in stressful environments (Furuta S et al., *Sweetpotato Res Front* (KNAES, Japan) 1:3, 1995; Sherwin HW and Farrant JM., *Plant Growth Regulation* 24: 203-210, 1998; and Yamasaki H *Trends in Plant Science* 2: 7-8, 1997).

Anthocyanins have demonstrated anti-oxidant activity, suggesting a role in protecting against cancer, cardiovascular and liver diseases (Kamei H et al., *J Clin Exp Med* 164: 829, 1993; Suda I, et al., 1997. *Sweetpotato Res Front* (KNAES, Japan) 4:3, 1997; and Wang CJ, et al., *J Food Chem Toxicology* 38: 411-416, 2000). Thus, anthocyanin-rich foods and extracts have been studied for their utility in a variety of therapeutic applications (e.g. Katsube et al., *J Agric Food Chem* (2003) 51(1):68-75; Renaud et al., *Lancet* (1992) 339:1523-1526; and Natella et al., *J Agric Food Chem* (2002)

50(26):7720-7725). There is also interest in the use of anthocyanin-rich plant species in the production of natural dyes (Venturi and Piccaglia, "*The Rediscovery of Dye Plants as Promising 'Non Food Crops'*", Interactive European Network for Industrial Crops and their Applications, Newsletter no. 10, November 1999).

Many steps in anthocyanin biosynthesis are shared among plant species, while the regulatory elements that underlie the expression level and pattern of genes encoding these enzymes are diverse. In *Petunia*, AN2 encodes a MYB domain protein that is orthologous to C1 from maize (Quattrocchio F et al., 1999, *Plant Cell* 11:1433-1444), and *Arabidopsis* genes PAP1 and PAP2 (Borevitz et al., *Plant Cell*. 2000 Dec;12(12):2383-2394). The Anthocyanin1 gene (AN1) of *petunia* encodes a basic helix-loop-helix (bHLH) protein that activates the transcription of the structural anthocyanin gene Dihydroflavonol Reductases (DFR). The expression of AN1 is regulated by AN2 (Spelt et al., *Plant Cell*. 2000 Sep;12(9):1619-32). In *Arabidopsis*, two other transcription factors have been implicated in controlling the accumulation of flavonoids: the homeodomain protein Anthocyaninless2 (ANL2) is required for anthocyanin accumulation in subepidermal cells, while and the zinc finger protein, TT1, is involved in the accumulation of proanthocyanidin polymers in the seed coat (Kubo et al., *Plant Cell*. 1999 Jul;11(7):1217-26.; Sagasser et al., *Genes Dev*. 2002 Jan 1;16(1):138-49).

Isoflavones have also been widely studied for their potential therapeutic utility and health benefits (Hewitt and Singletary, *Cancer Lett* (2003) 192(2):133-143; Katz, *J Altern Complement Med* (2002) 8(6):813-821). Isoflavones play roles in plant pathogen response and in symbioses with rhizobial bacteria (Pueppke et al. 1998, *Plant Physiol* 117:599-608). They occur almost exclusively in soybeans and other legumes (Jung et al. 2000, *Nature Biotechnology* 18:208-212). Three principle isoflavone aglycones occur in soybean: daidzein, genistein and glycitein. Glycitein accounts for only about 10% of the total isoflavone content (Song et al 1999, *J Agric Food Chem* 47:1607-1610), but some research suggests glycitein is both more bioavailable (Song et al 1999, *J of Nutr*. 129:957-962) and more estrogenic (Songe et al 1999, *J Agric Food Chem*, *supra*) than daidzein and genistein.

## SUMMARY OF THE INVENTION

The invention is directed to a method of obtaining flavonoids that comprises obtaining a plant that overexpresses an *ANT1* gene compared to wild-type plants, and extracting a flavonoid from the plant. In one embodiment of the invention, the plant is a

transgenic plant that contains a transformation vector that causes the overexpression of *ANT1*. In another embodiment, the plant has been selectively bred to have an allele of or mutation in an endogenous *ANT1* gene that causes the overexpression of *ANT1* compared to plants lacking the allele or mutation.

In one embodiment, the plant is tomato and the flavonoid extracted is an anthocyanin selected from the group consisting of delphinidin 3-rutinoside-5-glucoside, delphinidin 3-(coumaroyl)rutinoside-5-glucoside, delphinidin 3-(caffeoyl)rutinoside-5-glucoside, petunidin 3-rutinoside-5-glucoside, petunidin 3-(coumaroyl)rutinoside-5-glucoside, petunidin 3-(caffeoyl)rutinoside-5-glucoside, malvidin 3-rutinoside-5-glucoside, malvidin 3-(coumaroyl)rutinoside-5-glucoside, and malvidin 3-(caffeoyl)rutinoside-5-glucoside. Alternatively, the flavonoid extracted is glycitein.

The invention is also directed to flavonoid-containing plant extracts obtained from plants that overexpress *ANT1*.

### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 presents the core chemical structure of the anthocyanins listed in Table 2 below.

Figures 2a and 2b present the predicted chemical structures of the anthocyanins isolated from tobacco that over-expresses the *ANT1* gene, specifically cyanidin-3-glucoside (Fig. 2a) and cyanidin-3-rutinoside (Fig. 2b).

### DETAILED DESCRIPTION OF THE INVENTION

#### I. Definitions.

Unless otherwise indicated, all technical and scientific terms used herein have the same meaning as they would to one skilled in the art of the present invention.

Practitioners are particularly directed to Sambrook *et al.* Molecular Cloning: A Laboratory Manual (Second Edition), Cold Spring Harbor Press, Plainview, N.Y., 1989; and Ausubel *et al.* Current Protocols in Molecular Biology, John Wiley & Sons, New York, N.Y., 1993, for definitions and terms of the art.

All publications cited herein are expressly incorporated herein by reference for the purpose of describing and disclosing compositions and methodologies that might be used in connection with the invention. All cited patents, patent publications, and sequence and other information in referenced websites are also incorporated by reference.

As used herein, the term "vector" refers to a nucleic acid construct designed for transfer between different host cells. An "expression vector" refers to a vector that has the ability to incorporate and express heterologous DNA fragments in a foreign cell. Many prokaryotic and eukaryotic expression vectors are commercially available. Selection of appropriate expression vectors is within the knowledge of those having skill in the art.

A "heterologous" nucleic acid construct or sequence has a portion of the sequence which is not native to the plant cell in which it is expressed. Heterologous, with respect to a control sequence refers to a control sequence (*i.e.* promoter or enhancer) that does not function in nature to regulate the same gene the expression of which it is currently regulating. Generally, heterologous nucleic acid sequences are not endogenous to the cell or part of the genome in which they are present, and have been added to the cell, by infection, transfection, microinjection, electroporation, or the like. A "heterologous" nucleic acid construct may contain a control sequence/DNA coding sequence combination that is the same as, or different from a control sequence/DNA coding sequence combination found in the native plant.

As used herein, the term "gene" means the segment of DNA involved in producing a polypeptide chain, which may or may not include regions preceding and following the coding region, *e.g.* 5' untranslated (5' UTR) or "leader" sequences and 3' UTR or "trailer" sequences, as well as intervening sequences (introns) between individual coding segments (exons).

As used herein, "percent (%) sequence identity" with respect to a subject sequence, or a specified portion of a subject sequence, is defined as the percentage of nucleotides or amino acids in the candidate derivative sequence identical with the nucleotides or amino acids in the subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1990) 215:403-410; [blast.wustl.edu/blast/README.html](http://blast.wustl.edu/blast/README.html) website) with all the search parameters set to default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A % identity value is determined by the number of matching identical nucleotides or amino acids divided by the sequence length for which the percent identity is being reported. "Percent (%) amino acid sequence similarity" is determined by doing the same calculation as for determining % amino acid sequence identity, but including

conservative amino acid substitutions in addition to identical amino acids in the computation.

The term "% homology" is used interchangeably herein with the term "% identity."

A nucleic acid sequence is considered to be "selectively hybridizable" to a reference nucleic acid sequence if the two sequences specifically hybridize to one another under moderate to high stringency hybridization and wash conditions. Hybridization conditions are based on the melting temperature ( $T_m$ ) of the nucleic acid binding complex or probe. For example, "maximum stringency" typically occurs at about  $T_m - 5^\circ\text{C}$  ( $5^\circ$  below the  $T_m$  of the probe); "high stringency" at about  $5-10^\circ$  below the  $T_m$ ; "intermediate stringency" at about  $10-20^\circ$  below the  $T_m$  of the probe; and "low stringency" at about  $20-25^\circ$  below the  $T_m$ . Functionally, maximum stringency conditions may be used to identify sequences having strict identity or near-strict identity with the hybridization probe; while high stringency conditions are used to identify sequences having about 80% or more sequence identity with the probe.

Moderate and high stringency hybridization conditions are well known in the art (see, for example, Sambrook, *et al*, *supra*, Chapters 9 and 11, and in Ausubel, F.M., *et al*, *supra*). An example of high stringency conditions includes hybridization at about  $42^\circ\text{C}$  in 50% formamide, 5X SSC, 5X Denhardt's solution, 0.5% SDS and 100  $\mu\text{g/ml}$  denatured carrier DNA followed by washing two times in 2X SSC and 0.5% SDS at room temperature and two additional times in 0.1X SSC and 0.5% SDS at  $42^\circ\text{C}$ .

As used herein, "recombinant" includes reference to a cell or vector, that has been modified by the introduction of a heterologous nucleic acid sequence or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found in identical form within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all as a result of deliberate human intervention.

As used herein, the terms "transformed", "stably transformed" or "transgenic" with reference to a plant cell means the plant cell has a non-native (heterologous) nucleic acid sequence integrated into its genome which is maintained through two or more generations.

As used herein, the term "expression" refers to the process by which a polypeptide is produced based on the nucleic acid sequence of a gene. The process includes both transcription and translation.

The term "introduced" in the context of inserting a nucleic acid sequence into a cell, means "transfection", or "transformation" or "transduction" and includes reference to

the incorporation of a nucleic acid sequence into a eukaryotic or prokaryotic cell where the nucleic acid sequence may be incorporated into the genome of the cell (for example, chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (for example, transfected mRNA).

As used herein, a "plant cell" refers to any cell derived from a plant, including cells from undifferentiated tissue (*e.g.*, callus) as well as plant seeds, pollen, progagules and embryos.

As used herein, the terms "native" and "wild-type" relative to a given plant trait or phenotype refers to the form in which that trait or phenotype is found in the same variety of plant in nature.

As used herein, the term "modified" regarding a plant trait, refers to a change in the phenotype of a transgenic plant relative to a non-transgenic plant, as it is found in nature.

As used herein, the term " $T_1$ " refers to the generation of plants from the seed of  $T_0$  plants. The  $T_1$  generation is the first set of transformed plants that can be selected by application of a selection agent, *e.g.*, an antibiotic or herbicide, for which the transgenic plant contains the corresponding resistance gene.

As used herein, the term " $T_2$ " refers to the generation of plants by self-fertilization of the flowers of  $T_1$  plants, previously selected as being transgenic.

As used herein, the term "plant part" includes any plant organ or tissue including, without limitation, seeds, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores. Plant cells can be obtained from any plant organ or tissue and cultures prepared therefrom. The class of plants which can be used in the methods of the present invention is generally as broad as the class of higher plants amenable to transformation techniques, including both monocotyledenous and dicotyledenous plants.

As used herein, "transgenic plant" includes reference to a plant that comprises within its genome a heterologous polynucleotide. Generally, the heterologous polynucleotide is stably integrated within the genome such that the polynucleotide is passed on to successive generations. The heterologous polynucleotide may be integrated into the genome alone or as part of a recombinant expression cassette. "Transgenic" is used herein to include any cell, cell line, callus, tissue, plant part or plant, the genotype of which has been altered by the presence of heterologous nucleic acid including those transgenics initially so altered as well as those created by sexual crosses or asexual propagation from the initial transgenic.

Thus a plant having within its cells a heterologous polynucleotide is referred to herein as a "transgenic plant". The heterologous polynucleotide can be either stably integrated into the genome, or can be extra-chromosomal. Preferably, the polynucleotide of the present invention is stably integrated into the genome such that the polynucleotide is passed on to successive generations. The polynucleotide is integrated into the genome alone or as part of a recombinant expression cassette. "Transgenic" is used herein to include any cell, cell line, callus, tissue, plant part or plant, the genotype of which has been altered by the presence of heterologous nucleic acids including those transgenics initially so altered as well as those created by sexual crosses or asexual reproduction of the initial transgenics.

A plant cell, tissue, organ, or plant into which the recombinant DNA constructs containing the expression constructs have been introduced is considered "transformed", "transfected", or "transgenic". A transgenic or transformed cell or plant also includes progeny of the cell or plant and progeny produced from a breeding program employing such a transgenic plant as a parent in a cross and exhibiting an altered phenotype resulting from the presence of a recombinant nucleic acid sequence. Hence, a plant of the invention will include any plant which has a cell containing a construct with introduced nucleic acid sequences, regardless of whether the sequence was introduced into the directly through transformation means or introduced by generational transfer from a progenitor cell which originally received the construct by direct transformation.

The terms "*Anthocyanin 1*" and "*ANT1*", as used herein encompass native *Anthocyanin 1* (*ANT1*) nucleic acid and amino acid sequences, homologues, variants and fragments thereof.

An "isolated" *ANT1* nucleic acid molecule is an *ANT1* nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the *ANT1* nucleic acid. An isolated *ANT1* nucleic acid molecule is other than in the form or setting in which it is found in nature. However, an isolated *ANT1* nucleic acid molecule includes *ANT1* nucleic acid molecules contained in cells that ordinarily express *ANT1* where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

As used herein, the term "mutant" with reference to a polynucleotide sequence or gene differs from the corresponding wild type polynucleotide sequence or gene either in terms of sequence or expression, where the difference contributes to a modified plant phenotype or trait. Relative to a plant or plant line, the term "mutant" refers to a plant or

plant line which has a modified plant phenotype or trait, where the modified phenotype or trait is associated with the modified expression of a wild type polynucleotide sequence or gene.

Generally, a "variant" polynucleotide sequence encodes a "variant" amino acid sequence which is altered by one or more amino acids from the reference polypeptide sequence. The variant polynucleotide sequence may encode a variant amino acid sequence having "conservative" or "non-conservative" substitutions. Variant polynucleotides may also encode variant amino acid sequences having amino acid insertions or deletions, or both.

As used herein, the term "phenotype" may be used interchangeably with the term "trait". The terms refer to a plant characteristic that is readily observable or measurable and results from the interaction of the genetic make-up of the plant with the environment in which it develops. Such a phenotype includes chemical changes in the plant make-up resulting from enhanced gene expression which may or may not result in morphological changes in the plant, but which are measurable using analytical techniques known to those of skill in the art.

## II. The Identified *ANT1* Phenotype and Gene.

The gene and phenotype of this invention were identified in a screen using activation tagging. Activation tagging is a process by which a heterologous nucleic acid construct comprising a nucleic acid control sequence, *e.g.* an enhancer, is inserted into a plant genome. The enhancer sequences act to enhance transcription of one or more native plant genes (Walden *et al.*, EMBO J. 13: 4729-36, 1994; Walden *et al.*, *Plant Mol. Biol.* 26: 1521-8, 1994; and Weigel D, *et al.*, *Plant Physiology*, 122:1003-1013, 2000).

Briefly, a large number of tomato (*Lycopersium esculentum*) cv. Micro-Tom plants were transformed with a modified form of the activation tagging vector pSKI015 (Weigel *et al*, *supra*), which comprises a T-DNA (*i.e.*, the sequence derived from the Ti plasmid of *Agrobacterium tumefaciens* that are transferred to a plant cell host during *Agrobacterium* infection), an enhancer element and a selectable marker gene. The construct, pAG3202, is further described in the Examples. Following random insertion of pAG3202 into the genome of transformed plants, the enhancer element can result in up-regulation genes in the vicinity of the T-DNA insertion, generally within 5-10 kilobase (kb) of the insertion. In the T<sub>1</sub> generation, plants were exposed to the selective agent in order to specifically recover those plants that expressed the selectable marker and therefore harbored insertions



of the activation-tagging vector. Transformed plants were observed for interesting phenotypes, which are generally identified at the T<sub>1</sub>, T<sub>2</sub> and/or T<sub>3</sub> generations. Genomic sequence surrounding the T-DNA insertion is analyzed in order to identify genes responsible for the interesting phenotypes. Genes responsible for causing such phenotypes are identified as attractive targets for manipulation for agriculture, food, ornamental plant, and/or pharmaceutical industries.

The present invention provides a modified leaf, flower or fruit color phenotype, identified in ACTTAG Mico-Tom lines that were observed at the callus stage as having purple color and purple shoots. Purple plants were derived from purple colored caulogenic callus in culture. The clonal plant lines (i.e., additional shoots originating from the same purple colored caulogenic callus or those multiplied from the first purple plant either in tissue culture or by cuttings in the greenhouse) were identified as having purple coloration on leaves, sepals and flowers. The plants were also observed to exhibit a modified fruit color described as a deeper red color relative to wild type Micro-Tom plants. The phenotype and associated gene have been designated *Anthocyanin 1* ("*ANT1*").

The invention also provides a newly identified and isolated nucleic acid sequence that was identified by analysis of the genomic DNA sequence surrounding the T-DNA insertion correlating with the *ANT1* phenotype. In particular, applicants have identified and characterized the open reading frame of the *ANT1* gene, which is specifically overexpressed in plants having the *ANT1* phenotype, and which is provided in SEQ ID NO:1. A detailed description of the isolation and characterization of *ANT1* is set forth in the Examples.

### III. Compositions of the Invention

#### A. *ANT1* Nucleic acids

The *ANT1* gene may be used in the development of transgenic plants having a desired phenotype. This may be accomplished using the native *ANT1* sequence, a variant *ANT1* sequence or a homologue or fragment thereof.

An *ANT1* nucleic acid sequence of this invention may be a DNA or RNA sequence, derived from genomic DNA, cDNA or mRNA. The nucleic acid sequence may be cloned, for example, by isolating genomic DNA from an appropriate source, and amplifying and cloning the sequence of interest using PCR. Alternatively, nucleic acid sequence may be synthesized, either completely or in part, especially where it is desirable to provide plant-preferred sequences. Thus, all or a portion of the desired structural gene

(that portion of the gene which encodes a polypeptide or protein) may be synthesized using codons preferred by a selected host.

The invention provides a polynucleotide comprising a nucleic acid sequence which encodes or is complementary to a sequence which encodes an *ANT1* polypeptide having the amino acid sequence presented in SEQ ID NO:2 and a polynucleotide sequence identical over its entire length to the *ANT1* nucleic acid sequence presented SEQ ID NO:1. The invention also provides the coding sequence for the mature *ANT1* polypeptide, a variant or fragment thereof, as well as the coding sequence for the mature polypeptide or a fragment thereof in a reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, pro-, or prepro- protein sequence.

An *ANT1* polynucleotide can also include non-coding sequences, including for example, but not limited to, non-coding 5' and 3' sequences, such as the transcribed, untranslated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns, polyadenylation signals, and additional coding sequence that encodes additional amino acids. For example, a marker sequence can be included to facilitate the purification of the fused polypeptide. Polynucleotides of the present invention also include polynucleotides comprising a structural gene and the naturally associated sequences that control gene expression.

When an isolated polynucleotide of the invention comprises an *ANT1* nucleic acid sequence flanked by non- *ANT1* nucleic acid sequence, the total length of the combined polynucleotide is typically less than 25 kb, and usually less than 20kb, or 15 kb, and in some cases less than 10 kb, or 5 kb.

In addition to the *ANT1* nucleic acid and corresponding polypeptide sequences described herein, *ANT1* variants can be prepared by introducing appropriate nucleotide changes into the *ANT1* nucleic acid sequence; by synthesis of the desired *ANT1* polypeptide or by altering the expression level of the *ANT1* gene in plants. For example, amino acid changes may alter post-translational processing of the *ANT1* polypeptide, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

In one aspect, preferred *ANT1* coding sequences include a polynucleotide comprising a nucleic acid sequence which encodes or is complementary to a sequence which encodes an *ANT1* polypeptide having at least 50%, 60%, 70%, 75%, 80%, 85%, 90%, 95% or more sequence identity to the amino acid sequence presented in SEQ ID NO:2.

In another aspect, preferred variants include an *ANT1* polynucleotide sequence that is at least 50% to 60% identical over its entire length to the *ANT1* nucleic acid sequence presented as SEQ ID NO:1, and nucleic acid sequences that are complementary to such an *ANT1* sequence. More preferable are *ANT1* polynucleotide sequences comprise a region having at least 70%, 80%, 85%, 90% or 95% or more sequence identity to the *ANT1* sequence presented as SEQ ID NO:1.

In a related aspect, preferred variants include polynucleotides that are be "selectively hybridizable" to the *ANT1* polynucleotide sequence presented as SEQ ID NO:1.

Sequence variants also include nucleic acid molecules that encode the same polypeptide as encoded by the *ANT1* polynucleotide sequence described herein. Thus, where the coding frame of an identified nucleic acid molecule is known, for example by homology to known genes or by extension of the sequence, a number of coding sequences can be produced as a result of the degeneracy of the genetic code. For example, the triplet CGT encodes the amino acid arginine. Arginine is alternatively encoded by CGA, CGC, CGG, AGA, and AGG. Such substitutions in the coding region fall within the sequence variants that are covered by the present invention. Any and all of these sequence variants can be utilized in the same way as described herein for the identified *ANT1* parent sequence, SEQ ID NO:1.

Such sequence variants may or may not selectively hybridize to the parent sequence. This would be possible, for example, when the sequence variant includes a different codon for each of the amino acids encoded by the parent nucleotide. In accordance with the present invention, also encompassed are sequences that are at least 70% identical to such degeneracy-derived sequence variants.

Although *ANT1* nucleotide sequence variants are preferably capable of hybridizing to the nucleotide sequences recited herein under conditions of moderately high or high stringency, there are, in some situations, advantages to using variants based on the degeneracy of the code, as described above. For example, codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic organism, in accordance with the optimum codon usage dictated by the particular host organism. Alternatively, it may be desirable to produce RNA having longer half lives than the mRNA produced by the recited sequences.

Variations in the native full-length *ANT1* nucleic acid sequence described herein, may be made, for example, using any of the techniques and guidelines for conservative

and non-conservative mutations, as generally known in the art, oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis (Kunkel TA *et al.*, *Methods Enzymol.* 204:125-39, 1991); cassette mutagenesis (Cramer A and Stemmer WP, *Bio Techniques* 18(2):194-6, 1995.); restriction selection mutagenesis (Haught C *et al.* *BioTechniques* 16(1):47-48, 1994), or other known techniques can be performed on the cloned DNA to produce nucleic acid sequences encoding *ANT1* variants.

In addition, the gene sequences associated the *ANT1* phenotype may be synthesized, either completely or in part, especially where it is desirable to provide host-preferred sequences. Thus, all or a portion of the desired structural gene (that portion of the gene which encodes the protein) may be synthesized using codons preferred by a selected host. Host-preferred codons may be determined, for example, from the codons used most frequently in the proteins expressed in a desired host species.

It is preferred that an *ANT1* polynucleotide encodes an *ANT1* polypeptide that retains substantially the same biological function or activity as the mature *ANT1* polypeptide encoded by the polynucleotide set forth as SEQ ID NO:1 (i.e. results in an *ANT1* phenotype when overexpressed in a plant).

Variants also include fragments of the *ANT1* polynucleotide of the invention, which can be used to synthesize a full-length *ANT1* polynucleotide. Preferred embodiments include polynucleotides encoding polypeptide variants wherein 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues of an *ANT1* polypeptide sequence of the invention are substituted, added or deleted, in any combination. Particularly preferred are substitutions, additions, and deletions that are silent such that they do not alter the properties or activities of the polynucleotide or polypeptide.

A nucleotide sequence encoding an *ANT1* polypeptide can also be used to construct hybridization probes for further genetic analysis. Screening of a cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook *et al.*, *supra*). Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook *et al.*, *supra*.

The probes or portions thereof may also be employed in PCR techniques to generate a pool of sequences for identification of closely related *ANT1* sequences. When *ANT1* sequences are intended for use as probes, a particular portion of an *ANT1* encoding sequence, for example a highly conserved portion of the coding sequence may be used.

For example, an *ANT1* nucleotide sequence may be used as a hybridization probe for a cDNA library to isolate genes, for example, those encoding naturally-occurring variants of *ANT1* from other plant species, which have a desired level of sequence identity to the *ANT1* nucleotide sequence disclosed in SEQ ID NO:1. Exemplary probes have a length of about 20 to about 50 bases.

In another exemplary approach, a nucleic acid encoding an *ANT1* polypeptide may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein, and, if necessary, using conventional primer extension procedures as described in Sambrook *et al.*, *supra*, to detect *ANT1* precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

As discussed above, nucleic acid sequences of this invention may include genomic, cDNA or mRNA sequence. By "encoding" is meant that the sequence corresponds to a particular amino acid sequence either in a sense or anti-sense orientation. By "extrachromosomal" is meant that the sequence is outside of the plant genome of which it is naturally associated. By "recombinant" is meant that the sequence contains a genetically engineered modification through manipulation via mutagenesis, restriction enzymes, and the like.

Once the desired form of an *ANT1* nucleic acid sequence, homologue, variant or fragment thereof, is obtained, it may be modified in a variety of ways. Where the sequence involves non-coding flanking regions, the flanking regions may be subjected to resection, mutagenesis, etc. Thus, transitions, transversions, deletions, and insertions may be performed on the naturally occurring sequence.

With or without such modification, the desired form of the *ANT1* nucleic acid sequence, homologue, variant or fragment thereof, may be incorporated into a plant expression vector for transformation of plant cells.

#### B. *ANT1* Polypeptides

In one preferred embodiment, the invention provides an *ANT1* polypeptide, having a native mature or full-length *ANT1* polypeptide sequence comprising the sequence presented in SEQ ID NO:2. An *ANT1* polypeptide of the invention can be the mature *ANT1* polypeptide, part of a fusion protein or a fragment or variant of the *ANT1* polypeptide sequence presented in SEQ ID NO:2.

Ordinarily, an *ANTI* polypeptide of the invention has at least 50% to 60% identity to an *ANTI* amino acid sequence over its entire length. More preferable are *ANTI* polypeptide sequences that comprise a region having at least 70%, 80%, 85%, 90% or 95% or more sequence identity to the *ANTI* polypeptide sequence of SEQ ID NO:2.

Fragments and variants of the *ANTI* polypeptide sequence of SEQ ID NO:2, are also considered to be a part of the invention. A fragment is a variant polypeptide that has an amino acid sequence that is entirely the same as part but not all of the amino acid sequence of the previously described polypeptides. Exemplary fragments comprises at least 10, 20, 30, 40, 50, 75, or 100 contiguous amino acids of SEQ ID NO:2. The fragments can be "free-standing" or comprised within a larger polypeptide of which the fragment forms a part or a region, most preferably as a single continuous region. Preferred fragments are biologically active fragments, which are those fragments that mediate activities of the polypeptides of the invention, including those with similar activity or improved activity or with a decreased activity. Also included are those fragments that antigenic or immunogenic in an animal, particularly a human.

*ANTI* polypeptides of the invention also include polypeptides that vary from the *ANTI* polypeptide sequence of SEQ ID NO:2. These variants may be substitutional, insertional or deletional variants. The variants typically exhibit the same qualitative biological activity as the naturally occurring analogue, although variants can also be selected which have modified characteristics as further described below.

A "substitution" results from the replacement of one or more nucleotides or amino acids by different nucleotides or amino acids, respectively.

An "insertion" or "addition" is that change in a nucleotide or amino acid sequence which has resulted in the addition of one or more nucleotides or amino acid residues, respectively, as compared to the naturally occurring sequence.

A "deletion" is defined as a change in either nucleotide or amino acid sequence in which one or more nucleotides or amino acid residues, respectively, are absent.

Amino acid substitutions are typically of single residues; insertions usually will be on the order of from about 1 to 20 amino acids, although considerably larger insertions may be tolerated. Deletions range from about 1 to about 20 residues, although in some cases deletions may be much larger.

Substitutions, deletions, insertions or any combination thereof may be used to arrive at a final derivative. Generally these changes are done on a few amino acids to minimize the

alteration of the molecule. However, larger changes may be tolerated in certain circumstances.

Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, *i.e.*, conservative amino acid replacements. Insertions or deletions may optionally be in the range of 1 to 5 amino acids.

Substitutions are generally made in accordance with known "conservative substitutions". A "conservative substitution" refers to the substitution of an amino acid in one class by an amino acid in the same class, where a class is defined by common physicochemical amino acid side chain properties and high substitution frequencies in homologous proteins found in nature (as determined, *e.g.*, by a standard Dayhoff frequency exchange matrix or BLOSUM matrix). (*See generally, Doolittle, R.F., OF URFS and ORFS* (University Science Books, CA, 1986.))

A "non-conservative substitution" refers to the substitution of an amino acid in one class with an amino acid from another class.

*ANT1* polypeptide variants typically exhibit the same qualitative biological activity as the naturally occurring analogue, although variants also are selected to modify the characteristics of the *ANT1* polypeptide, as needed. For example, glycosylation sites, and more particularly one or more O-linked or N-linked glycosylation sites may be altered or removed. For example, amino acid changes may alter post-translational processes of the *ANT1* polypeptide, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis (Carter *et al.*, *Nucl. Acids Res.* **13**:4331, 1986; Zoller *et al.*, *Nucl. Acids Res.* **10**:6487, 1987), cassette mutagenesis (Wells *et al.*, *Gene* **34**:315, 1985), restriction selection mutagenesis (Wells *et al.*, *Philos. Trans. R. Soc. London SerA* **317**:415, 1986) or other known techniques can be performed on the cloned DNA to produce the *ANT1* polypeptide-encoding variant DNA.

Also included within the definition of *ANT1* polypeptides are other related *ANT1* polypeptides. Thus, probe or degenerate PCR primer sequences may be used to find other related polypeptides. Useful probe or primer sequences may be designed to all or part of the *ANT1* polypeptide sequence, or to sequences outside the coding region. As is generally known in the art, preferred PCR primers are from about 15 to about 35 nucleotides in length,

with from about 20 to about 30 being preferred, and may contain inosine as needed. The conditions for the PCR reaction are generally known in the art.

Covalent modifications of *ANTI* polypeptides are also included within the scope of this invention. For example, the invention provides *ANTI* polypeptides that are a mature protein and may comprise additional amino or carboxyl-terminal amino acids, or amino acids within the mature polypeptide (for example, when the mature form of the protein has more than one polypeptide chain). Such sequences can, for example, play a role in the processing of a protein from a precursor to a mature form, allow protein transport, shorten or lengthen protein half-life, or facilitate manipulation of the protein in assays or production. Cellular enzymes can be used to remove any additional amino acids from the mature protein (Creighton, T.E., *PROTEINS: STRUCTURE AND MOLECULAR PROPERTIES*, W.H. Freeman & Co., San Francisco, pp. 79-86, 1983).

In a preferred embodiment, overexpression of an *ANTI* polypeptide or variant thereof is associated with the *ANTI* phenotype.

### C. Antibodies.

The present invention further provides anti-*ANTI* polypeptide antibodies. The antibodies may be polyclonal, monoclonal, humanized, bispecific or heteroconjugate antibodies.

Polyclonal antibodies can be produced in a mammal, for example, following one or more injections of an immunizing agent, and preferably, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected into the mammal by a series of subcutaneous or intraperitoneal injections. The immunizing agent may include an *ANTI* polypeptide or a fusion protein thereof. It may be useful to conjugate the antigen to a protein known to be immunogenic in the mammal being immunized.

Alternatively, the anti-*ANTI* polypeptide antibodies may be monoclonal antibodies. Monoclonal antibodies may be produced by hybridomas, wherein a mouse, hamster, or other appropriate host animal, is immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent (Kohler and Milstein, *Nature* 256:495, 1975). Monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567.

The anti-*ANTI* polypeptide antibodies of the invention may further comprise humanized antibodies or human antibodies. The term "humanized antibody" refers to humanized forms of non-human (*e.g.*, murine) antibodies that are chimeric antibodies,



immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub> or other antigen-binding partial sequences of antibodies) which contain some portion of the sequence derived from non-human antibody. Methods for humanizing non-human antibodies are well known in the art, as further detailed in Jones *et al.*, *Nature* 321:522-525, 1986; Riechmann *et al.*, *Nature* 332:323-327, 1988; and Verhoeyen *et al.*, *Science* 239:1534-1536, 1988. Methods for producing human antibodies are also known in the art. (Jakobovits, A, *et al.*, *Ann N Y Acad Sci* 764:525-35, 1995; Jakobovits, A, *Curr Opin Biotechnol*, 6(5):561-6, 1995.

In one exemplary approach, anti-*ANT1* polyclonal antibodies are used for gene isolation. Western blot analysis may be conducted to determine that *ANT1* or a related protein is present in a crude extract of a particular plant species. When reactivity is observed, genes encoding the related protein may be isolated by screening expression libraries representing the particular plant species. Expression libraries can be constructed in a variety of commercially available vectors, including lambda gt11, as described in Sambrook, *et al.*, *supra*.

#### IV. Utility Of the *ANT1* Phenotype and Gene

From the foregoing, it can be appreciated that the *ANT1* nucleotide sequence, protein sequence and phenotype find utility in modulated expression of the *ANT1* protein and the development of non-native phenotypes associated with such modulated expression.

The *ANT1* phenotype has features that distinguish the mutant from wild type plants, including modified leaf color, modified flower color and modified fruit color. We have shown that the modified pigmentation phenotype is associated with increased production of specific anthocyanins, which vary according to individual plant species.

In one aspect, the modified leaf, flower and fruit color of plants having the *ANT1* phenotype finds utility in the development of improved ornamental plants, fruits and/or cut flowers.

In another aspect, the modified anthocyanin content in plants having the *ANT1* phenotype finds utility in plant-derived food, food additives, nutrition supplements, and natural dyes.

The *ANT1* gene may be used to generate transgenic plants that produce flavonoids including anthocyanins and isoflavones. When separation from other plant material is desired, flavonoids may be extracted by any method known in the art (Yang *et al.*, *J Chromatogr A* (2001) 928(2):163-170; Di Mauro *et al.*, *J. Agric. Food Chem* (2002)

50:5968-5974; Matsumoto et al., J. Agric. Food Chem (2001) 49:1541-1545). An extracted flavonoid may be substantially purified or may be used in an unprocessed or partially processed state.

In one preferred embodiment, the invention provides transgenic tomato that produces at least one anthocyanin selected from delphinidin 3-rutinoside-5-glucoside, delphinidin 3-(coumaroyl)rutinoside-5-glucoside, delphinidin 3-(caffeoyl)rutinoside-5-glucoside, petunidin 3-rutinoside-5-glucoside, petunidin 3-(coumaroyl)rutinoside-5-glucoside, petunidin 3-(caffeoyl)rutinoside-5-glucoside, malvidin 3-rutinoside-5-glucoside, malvidin 3-(coumaroyl)rutinoside-5-glucoside, and malvidin 3-(caffeoyl)rutinoside-5-glucoside. In a further preferred embodiment, the anthocyanin is produced at a level that is at least 5-, 10-, 20-, 50-, or 100-fold that observed in the non-transgenic plant.

In another preferred embodiment, the invention provides transgenic tobacco that produces at least one anthocyanin selected from cyanidin-3-glucoside and cyanidin-3-rutinoside. In a further preferred embodiment, the anthocyanin is produced at a level that is at least 5-, 10-, 20-, 50-, or 100-fold that observed in the non-transgenic plant.

We have further found that over-expression of the *ANT1* gene in tomato results in isoflavone production, which is otherwise undetectable. Accordingly, *ANT1* genes can be used in the generation of transgenic soy or other legumes with altered isoflavone content or composition. *ANT1* genes can also be used to produce isoflavones in plants other than legumes. In one embodiment, plants are generated that have increased glycitein content. In another embodiment, the isoflavone is produced at a level of at least 1.00 mg/100g. Thus, the *ANT1* gene may be used to generate transgenic plants that produce desired metabolites, including isoflavones. The isoflavones may be extracted by any method known in the art.

In another aspect, as further described in the Examples, the *ANT1* gene has utility as a transformation marker in genetically manipulated plants.

In practicing the invention, the *ANT1* phenotype and modified *ANT1* expression is generally applicable to any type of plant, as further detailed below.

The methods described herein are generally applicable to all plants. Although activation tagging and gene identification was carried out in tomato, following identification of a nucleic acid sequence and associated phenotype, the selected gene, a homologue, variant or fragment thereof, may be expressed in any type of plant. In one aspect, the invention is directed to fruit- and vegetable-bearing plants.

The invention is generally applicable to plants which produce fleshy fruits; for example but not limited to, tomato (*Lycopersicum*); grape (*Vitis*); strawberry (*Fragaria*); raspberry, blackberry, loganberry (*Rubus*); currants and gooseberry (*Ribes*); blueberry, bilberry, whortleberry, cranberry (*Vaccinium*); kiwifruit and Chinese gooseberry (*Actinidia*); apple (*Malus*); pear (*Pyrus*); melons (*Cucumis sp.*) members of the *Prunus* genera, e.g. plum, cherry, nectarine and peach; sapota (*Manilkara zapotilla*); mango; avocado; apricot; peaches; cherries; pineapple; papaya; passion fruit; citrus; date palm; banana; plantain; and fig.

Similarly, the invention is applicable to vegetable plants, including, but not limited to sugar beets, green beans, broccoli, brussels sprouts, cabbage, celery, chard, cucumbers, eggplants, peppers, pumpkins, rhubarb, winter squash, summer squash, zucchini, lettuce, radish, carrot, pea, potato, corn, murraya and herbs.

In a related aspect, the invention is directed to the cut flower industry, grain-producing plants, oil-producing plants and nut-producing plants, as well as other crops including, but not limited to, cotton (*Gossypium*), alfalfa (*Medicago sativa*), flax (*Linum usitatissimum*), tobacco (*Nicotiana*), turfgrass (*Poaceae* family), and other forage crops. Suitable transformation techniques for these and other plants are known in the art.

A wide variety of transformation techniques exist in the art, and new techniques are continually becoming available. Any technique that is suitable for the target host plant can be employed within the scope of the present invention. For example, the constructs can be introduced in a variety of forms including, but not limited to as a strand of DNA, in a plasmid, or in an artificial chromosome. The introduction of the constructs into the target plant cells can be accomplished by a variety of techniques, including, but not limited to *Agrobacterium*-mediated transformation, electroporation, microinjection, microprojectile bombardment calcium-phosphate-DNA co-precipitation or liposome-mediated transformation of a heterologous nucleic acid construct comprising the *ANT1* coding sequence. The transformation of the plant is preferably permanent, *i.e.* by integration of the introduced expression constructs into the host plant genome, so that the introduced constructs are passed onto successive plant generations.

In one embodiment, binary Ti-based vector systems may be used to transfer and confirm the association between enhanced expression of an identified gene with a particular plant trait or phenotype. Standard *Agrobacterium* binary vectors are known to those of skill in the art and many are commercially available, such as pBI121 (Clontech Laboratories, Palo Alto, CA).

The optimal procedure for transformation of plants with *Agrobacterium* vectors will vary with the type of plant being transformed. Exemplary methods for *Agrobacterium*-mediated transformation include transformation of explants of hypocotyl, shoot tip, stem or leaf tissue, derived from sterile seedlings and/or plantlets. Such transformed plants may be reproduced sexually, or by cell or tissue culture. *Agrobacterium* transformation has been previously described for a large number of different types of plants and methods for such transformation may be found in the scientific literature.

Depending upon the intended use, a heterologous nucleic acid construct may be made which comprises a nucleic acid sequence associated with the *ANT1* phenotype, and which encodes the entire protein, or a biologically active portion thereof for transformation of plant cells and generation of transgenic plants.

The expression of an *ANT1* nucleic acid sequence or a homologue, variant or fragment thereof may be carried out under the control of a constitutive, inducible or regulatable promoter. In some cases expression of the *ANT1* nucleic acid sequence or homologue, variant or fragment thereof may be regulated in a developmental stage or tissue-associated or tissue-specific manner. Accordingly, expression of the nucleic acid coding sequences described herein may be regulated with respect to the level of expression, the tissue type(s) where expression takes place and/or developmental stage of expression leading to a wide spectrum of applications wherein the expression of an *ANT1* coding sequence is modulated in a plant.

Strong promoters with enhancers may result in a high level of expression. When a low level of basal activity is desired, a weak promoter may be a better choice. Expression of *ANT1* nucleic acid sequence or homologue, variant or fragment thereof may also be controlled at the level of transcription, by the use of cell type specific promoters or promoter elements in the plant expression vector.

Numerous promoters useful for heterologous gene expression are available. Exemplary constitutive promoters include the raspberry E4 promoter (U.S. Patent Nos. 5,783,393 and 5,783,394), the 35S CaMV (Jones JD *et al.*, Transgenic Res 1:285-297 1992), the CsVMV promoter (Verdaguer B *et al.*, Plant Mol Biol 37:1055-1067, 1998) and the melon actin promoter. Exemplary tissue-specific promoters include the tomato E4 and E8 promoters (U.S. Patent No. 5,859,330) and the tomato 2AII gene promoter (Van Haaren MJJ *et al.*, Plant Mol Bio 21:625-640, 1993).

When *ANT1* sequences are intended for use as probes, a particular portion of an *ANT1* encoding sequence, for example a highly conserved portion of a coding sequence may be used.

In yet another aspect, in some cases it may be desirable to inhibit the expression of endogenous *ANT1* sequences in a host cell. Exemplary methods for practicing this aspect of the invention include, but are not limited to antisense suppression (Smith, *et al.*, *Nature* 334:724-726, 1988); co-suppression (Napoli, *et al.*, *Plant Cell* 2:279-289, 1990); ribozymes (PCT Publication WO 97/10328); and combinations of sense and antisense (Waterhouse, *et al.*, *Proc. Natl. Acad. Sci. USA* 95:13959-13964, 1998). Methods for the suppression of endogenous sequences in a host cell typically employ the transcription or transcription and translation of at least a portion of the sequence to be suppressed. Such sequences may be homologous to coding as well as non-coding regions of the endogenous sequence. In some cases, it may be desirable to inhibit expression of the *ANT1* nucleotide sequence. This may be accomplished using procedures generally employed by those of skill in the art together with the *ANT1* nucleotide sequence provided herein.

Standard molecular and genetic tests may be performed to analyze the association between a cloned gene and an observed phenotype. A number of other techniques that are useful for determining (predicting or confirming) the function of a gene or gene product in plants are described below.

#### 1. DNA/RNA analysis

DNA taken from a mutant plant may be sequenced to identify the mutation at the nucleotide level. The mutant phenotype may be rescued by overexpressing the wild type (WT) gene. The stage- and tissue-specific gene expression patterns in mutant vs. WT lines, for instance, by in situ hybridization, may be determined. Analysis of the methylation status of the gene, especially flanking regulatory regions, may be performed. Other suitable techniques include overexpression, ectopic expression, expression in other plant species and gene knock-out (reverse genetics, targeted knock-out, viral induced gene silencing (Baulcombe D, *Arch Virol Suppl* 15:189-201, 1999).

In a preferred application, microarray analysis, also known as expression profiling or transcript profiling, is used to simultaneously measure differences or induced changes in the expression of many different genes. Techniques for microarray analysis are well known in the art (Schena M *et al.*, *Science* (1995) 270:467-470; Baldwin D *et al.*, *Curr Opin Plant Biol.* 2(2):96-103, 1999; Dangond F, *Physiol Genomics* (2000) 2:53-58; van Hal NL *et al.*, *J Biotechnol* (2000) 78:271-280; Richmond T and Somerville S, *Curr Opin*

Plant Biol (2000) 3:108-116). Microarray analysis of individual tagged lines may be carried out, especially those from which genes have been isolated. Such analysis can identify other genes that are coordinately regulated as a consequence of the overexpression of the gene of interest, which may help to place an unknown gene in a particular pathway.

## **2. Gene Product Analysis**

Analysis of gene products may include recombinant protein expression, antisera production, immunolocalization, biochemical assays for catalytic or other activity, analysis of phosphorylation status, and analysis of interaction with other proteins via yeast two-hybrid assays.

## **3. Pathway Analysis**

Pathway analysis may include placing a gene or gene product within a particular biochemical or signaling pathway based on its overexpression phenotype or by sequence homology with related genes. Alternatively, analysis may comprise genetic crosses with WT lines and other mutant lines (creating double mutants) to order the gene in a pathway, or determining the effect of a mutation on expression of downstream “reporter” genes in a pathway.

## **4. Other Analyses**

Other analyses may be performed to determine or confirm the participation of the isolated gene and its product in a particular metabolic or signaling pathway, and to help determine gene function.

## **Generation of Mutated Plants with an ANT1 Phenotype**

The invention further provides a method of identifying plants that have mutations in, or an allele of, endogenous *ANT1* that confer an *ANT1* phenotype, and generating progeny of these plants that also have the *ANT1* phenotype and are not genetically modified. In one method, called “TILLING” (for Targeting Induced Local Lesions IN Genomes), mutations are induced in the seed of a plant of interest, for example, using EMS treatment. The resulting plants are grown and self-fertilized, and the progeny are used to prepare DNA samples. *ANT1*-specific PCR is used to identify whether a mutated plant has an *ANT1* mutation. Plants having *ANT1* mutations may then be tested for the *ANT1* phenotype, or alternatively, plants may be tested for the *ANT1* phenotype, and then *ANT1*-specific PCR is used to determine whether a plant having the *ANT1* phenotype has a mutated *ANT1* gene. TILLING can identify mutations that may alter the expression of

specific genes or the activity of proteins encoded by these genes (see Colbert et al (2001) Plant Physiol 126:480-484; McCallum et al (2000) Nature Biotechnology 18:455-457).

In another method, a candidate gene/Quantitative Trait Locus (QTLs) approach can be used in a marker-assisted breeding program to identify alleles of or mutations in the *ANT1* gene or orthologs of *ANT1* that may confer the *ANT1* phenotype (see Foolad et al., Theor Appl Genet. (2002) 104(6-7):945-958; Rothan et al., Theor Appl Genet (2002) 105(1):145-159); Dekkers and Hospital, Nat Rev Genet. (2002) Jan;3(1):22-32).

Thus, in a further aspect of the invention, an *ANT1* nucleic acid is used to identify whether a plant having an *ANT1* phenotype has a mutation in endogenous *ANT1* or has a particular allele that causes the *ANT1* phenotype compared to plants lacking the mutation or allele, and generating progeny of the identified plant that have inherited the *ANT1* mutation or allele and have the *ANT1* phenotype. The *ANT1* plants generated can be used as non-genetically modified foods having increased flavonoid content, and can also be used for the same purposes described herein for transgenic *ANT1* plants (e.g. extraction of natural dyes, etc.).

### EXAMPLE 1

#### Generation of Plants with an *ANT1* Phenotype by Transformation with an Activation Tagging Construct

##### A. *Agrobacterium* vector preparation.

Mutants were generated using a modified version of the activation tagging "ACTTAG" vector, pSKI015 (GenBank Identifier [GI] 6537289; Weigel D *et al.*, *supra*). This binary vector, called pAG3202, contains the following components: the pSKI backbone; a 4X 35S enhancer consisting of four tandem repeats of the enhancer region from the CaMV 35S promoter including 4 Alu1-EcoRV fragments in tandem, 129 bp of CaMV sequence associated with each tandem Alu1-EcoRV repeat, and an additional 7 bp repeated sequence that is not in the 35S enhancer region of the native CaMV genome; the *np1II* selectable marker under the control of a raspberry E4 (RE4) promoter; an *Agrobacterium* gene 7 termination element located downstream of the *np1II* gene, adjacent the left border of the plasmid.

Single colonies of *Agrobacterium tumefaciens* strains EHA 105/EHA 101/GV3101 containing the binary plasmid pAG3202 were grown in MGL medium at pH 5.4 overnight and diluted to approximately  $5 \times 10^8$  cells/ml with MGL or liquid plant co-cultivation medium.

For long-term storage, PCR-positive colonies were grown in selective media, glycerol added to a final concentration of 30% and cultures quick frozen, then stored at -80°C. For the initiation of dense *Agrobacterium* cultures for plant transformation, stock cultures were grown in selective media, glycerol added to a final concentration of 30%, and a number of 20 µl aliquots quick frozen in liquid nitrogen and stored at -80°C.

#### **B. Transformation and Selection of *Micro-Tom* Mutants**

Seeds of (*Lycopersium esculentum*) were surface sterilized in 25% bleach with tween-20 for 15 minutes and rinsed with sterile water before plating on seed germination medium (MS salts, Nitsch vitamins, 3% sucrose and 0.7% agar, pH 5.8), modified by the addition of auxin and/or cytokinins and giberellic acid as necessary. The cultures were incubated at 24°C with a 16 hr photo period (50-60 µmol.m<sup>-2</sup>s<sup>-1</sup>). Seven to ten day old seedlings and one month old *in vitro* plants were used for hypocotyl explants.

Hypocotyls were cut into 3-5 mm segments, then immersed in bacterial suspension, blotted on sterile filter paper and placed on co-cultivation medium. The explants were immersed in bacterial suspension, blotted on sterile filter paper and placed on co-cultivation medium (MS salts, LS vitamins, 3% sucrose, 0.1 mg/l kinetin, 0.2 mg/l 2,4-D, 200 mg/l potassium acid phosphate, 50 µM acetosyringone and 0.7% agar, pH 5.4) for 2-3 days.

After two to three days of co-culture, the explants were transferred to shoot regeneration medium containing MS salts, Nitsch vitamins, 3% sucrose, 2 mg/l zeatin, 500 mg/l carbenicillin, 200mg/L timetin and 0.7% agar at pH 5.8, supplemented with the antibiotic, kanamycin at 75 – 400 mg/l in order to select for *nptII*-expressing transformants. The selection level of antibiotic was gradually raised over an 8-week period based on the tissue response.

The explants were transferred to fresh medium every two weeks. Initiation of callus with signs of shoot initials was observed from 3-6 weeks depending on the type of explant. Callusing and shoot regeneration was observed to continue over approximately 4 months after which the explant tissues decline. A purple callus was observed among the tissue growing on the selection medium. Regenerated shoots displayed a variety of color phenotypes and were entirely green, entirely purple, or mix of green and purple to various degrees. Green shoots of approximately 1 cm in size with distinct shoot meristems were excised from the callus and transferred to root induction medium containing MS salts, Nitsch vitamins, 3% sucrose, 1 mg/l IBA, 50 mg/l kanamycin, 100 mg/l carbenicillin or



100mg/L timetin and 0.7% agar, pH 5.8. The rooted plants were out-planted to soil in a Biosafety greenhouse.

Plants were transported to greenhouse facilities, potted up in 3.5" pots tagged for plant identification.

Transformants were observed at the callus stage and after T<sub>1</sub> plants were established in the greenhouse for phenotypic variations relative to wild-type Micro-Tom plants. To achieve this, several wild-type plants were kept in close proximity to the transgenic plants. Each plant was observed closely twice a week with observations noted and documented by photographs.

Images of each pool of 8 plants were recorded using a Digital camera (DC-260), and morphology observations were made at about four weeks after planting.

Eleven Micro-Tom lines were developed from the callus originally identified by its purple color and purple shoots at the caulogenic stage. The clonal plant lines were identified as having modified leaf color with a heavy purple cast on leaves, modified flower color characterized by purple striations on petals and sepals and flowers with a purple cast mixed with the normal yellow color of the corolla. The plants were also observed to exhibit a modified fruit color described as a deeper red color relative to wild type Micro-Tom plants. The clonal plant lines (mutants) were designated *Anthocyanin1* ("ANT1").

The *ANT1* mutant was identified from fewer than 2000 individual Micro-Tom tomato ACTTAG lines that were developed following tissue culture transformation with the binary plasmid pAG3202, and selection on kanamycin-containing medium.

Observations were made and photos taken of the clonal T<sub>1</sub> *ANT1* plant lines that exhibited the *ANT1* phenotype, designated H000001484, H000001624, H000001708, H000001709, H000001710, H000001711, H000001712, H000001713, H000001715, H000001716 and H000001717.

Seeds were collected from T<sub>1</sub> plants from line H000001624 and grown to generate T<sub>2</sub> plants. From the 11 out of the 18 seeds that germinated, and 8 plants displayed purple coloration, confirming that *ANT1* is a dominant, gain of function mutation, following typical Mendelian segregation.

The results indicated that *ANT1* is a gain of function trait, expected from activation tagging based over-expression of a native gene.

## EXAMPLE 2

### Characterization of Plants That Exhibit the ANTI Phenotype.

Micro-Tom genomic DNA was extracted from the H000001484 clone of the activation tagged mutant originally identified at the callus stage, in sufficient yield and quality for plasmid rescue of activation tagged plant lines using the procedure described below. Further analysis was performed using combined tissue derived from the H000001624, H000001708, H000001709, H000001710, H000001711, H000001712, H000001713, H000001715, H000001716 and H000001717 plant lines.

#### A. Micro-Tom Tomato Genomic DNA Extraction

Nucleon™ PhytoPure™ systems (Plant and fungal DNA extraction kits) from Amersham™ were used for extracting genomic DNA. Methods were essentially as follows:

1.0g of fresh tissue from the H000001484 clone was ground in liquid nitrogen to yield a free flowing powder, then transferred to a 15 ml polypropylene centrifuge tube. 4.6 ml of Reagent 1 from the Nucleon Phyt PURE kit was added with thorough mixing, followed by addition of 1.5 ml of Reagent 2 from the Nucleon Phyt PURE kit, with inversion until a homogeneous mixture was obtained. The mixture was incubated at 65°C in a shaking water bath for 10 minutes, and placed on ice for 20 minutes. The samples were removed from the ice, 2 ml of – 20°C chloroform added, mixed and centrifuged at 1300g for 10 minutes. The supernatant was transferred into a fresh tube, 2 ml cold chloroform, 200 µl of Nucleon PhytoPure DNA extraction resin suspension added and the mixture shaken on a tilt shaker for 10 minutes at room temperature, then centrifuged at 1300g for 10 minutes. Without disturbing the Nucleon resin suspension layer, the upper DNA-containing phase was transferred into a fresh tube, centrifuged at 9500 rpm for 30 minutes to clarify the transferred aqueous phase if the upper phase appeared cloudy, an equal volume of cold isopropanol added, and the tube gently inverted until DNA precipitated. It was then pelleted by centrifugation, washed with cold 70% ethanol, pelleted again, and air-dried.

DNA was resuspended in TE buffer (10 mM Tris. HCl, pH 7.4, 1 mM EDTA), containing RNase, incubated at 55° C for 15 minutes, further extracted phenol/chloroform, then chloroform, run on a 1% agarose gel to check the DNA Quality, the DNA concentration determined by a DNA fluorometer (Hoeffer DyNA Quant 200).

DNA extracted from shoots of the H000001484 *ANT1* clone at the caulogenic callus stage and from wild type plants was PCR-amplified using primers that amplify a 35S enhancer sequence, and primers that amplify a region of the pBluescript vector sequence in pAG3202. Amplification using primers that span the 35S enhancer region resulted in a ladder of products, indicating that all four copies of the 35S enhancer were present. Amplification using primers to the pBluescript vector was done primarily to detect the T-DNA insert(s) in transformed plants and has been optimized for the following conditions: annealing temp: 57°C, 30 cycles [94°C, 30sec; 57°C, 1 min; 72°C, 1 min] 1 cycle [72°C, 7 min].

The ACTTAG™ line, H000001484 (*ANT1*), was confirmed as positive for the presence of 35S enhancer and pAG3202 vector sequences by PCR, and as positive for Southern hybridization verifying genomic integration of the ACTTAG DNA and showing the presence of a single T-DNA insertion in the clonal transgenic line.

#### B. Plasmid Rescue

Genomic DNA from the H000001484 clonal line was digested by the restriction enzymes used in Southern Hybridization. The restriction fragments were self-ligated and used to transform the *E. coli* cells. The plasmids that contained a full-length pBluescript vector, 4X 35S enhancer, and a right border T-DNA flanking genomic DNA fragment were rescued.

More specifically, genomic DNA was digested with Hind III and Xho I under standard reaction conditions at 37°C overnight.

The ligation reactions were set up containing the following and left at 16°C overnight:

Digested Genomic DNA	40 µl
5X Ligation Buffer	50 µl
Ligase (Gibcol, 1U/µl)	10 µl
ddH <sub>2</sub> O	150 µl

The ligated DNA precipitated, resuspended in ddH<sub>2</sub>O and used to transform *E. coli* SURE cells (Stratagene) *via* electroporation, with 10 pg of pUC18 plasmid as a control.

The transformation mixture was spread on two LB-plates containing 100 µg/ml ampicillin and incubated overnight at 37°C. Single colonies were picked from the plates and used to start a 5 ml LB-ampicillin broth culture from each colony by culturing

overnight at 37°C. The plasmid was extracted from the culture and restriction digested to confirm the size of genomic insertion.

### C. Sequencing Of Rescued Plasmids

Sequencing was accomplished using a ABI Prism BigDye™ Terminator Cycle Sequencing Ready Reaction Kit (PE Applied Biosystem), AmpliTaq DNA Polymerase (Perkin-Elmer), an ABI Prism™ 310 Genetic Analyzer (Perkin-Elmer) and sequence analysis software, e.g., Sequencer™ 3.1.1 or MacVector 6.5.3. Sequencing was done essentially according to manufacturers' protocols

The left ends of plasmids rescued were sequenced across the right T-DNA border.

The rescued sequence was subjected to analysis using the BLAST sequence comparison programs at the [www.ncbi.nlm.nih.gov/BLAST](http://www.ncbi.nlm.nih.gov/BLAST) website. A basic BLASTN search identified a sequence with 31% identity to the Anthocyanin 2 (An2) mRNA of *Petunia integrifolia* (GI 7673087 and 7673085). The presence of an open reading frame (i.e., the *ANT1* cDNA) was predicted using the BLASTX program.

RT-PCR analysis confirmed that the gene whose nucleotide sequence is presented as SEQ ID NO:1 (*ANT1*) was specifically overexpressed in tissue from plants having the *ANT1* phenotype. Specifically, RNA was extracted from combined tissues derived from the H000001624 clonal plant line, which exhibited the *ANT1* phenotype, and from wild type plants. RT-PCR was performed using primers specific to the sequence presented as SEQ ID NO:1 and a constitutively expressed actin gene (positive control). The results showed that plants displaying the *ANT1* phenotype over-expressed the mRNA for the *ANT1* gene, indicating the enhanced expression of the *ANT1* gene correlated with the *ANT1* phenotype.

The amino acid sequence predicted from the *ANT1* nucleic acid sequence was determined using Vector NTI (InforMax, North Bethesda, MD) and is presented in SEQ ID NO:2. A Basic BLASTP 2.0.11 search using the [ncbi.nlm.nih.gov/BLAST](http://ncbi.nlm.nih.gov/BLAST) website and the predicted *ANT1* amino acid sequence was conducted. Results indicated that the predicted *ANT1* protein sequence has 49% identity to the *Petunia integrifolia* An2 protein sequence (GI 7673088 and 7673086) and 65%-85% identity to several *Myb*-related transcription factors in the N-terminal region, from approximately aa 1-120 of SEQ ID NO:2. These *Myb*-related proteins included An2 from *Petunia x hybrida* (GI 7673084), the *Zea mays* C1-I (GI 22214), the *Zea mays* PL transcription factor (GI 2343273) and an

*Arabidopsis* transcription factor (GI 3941508). The *Petunia An2* gene is a regulator of the Anthocyanin biosynthetic pathway (Quattrocchio et al, *supra*).

These results suggest that *ANT1* is associated with modified leaf, flower or fruit color in Micro-tomato.

### EXAMPLE 3

#### Confirmation of Phenotype/Genotype Association in Micro-tomato

In order to further confirm the association between the *ANT1* phenotype and the *ANT1* gene presented in SEQ ID NO: 1, a genomic fragment comprising the *ANT1* gene, provided in SEQ ID NO:3, was over-expressed in wild type Micro-Tom plants. Specifically, this 1012 bp genomic fragment, including the *ANT1* coding regions, was cloned into the multiple cloning site (MCS) of the binary vector pAG2370. pAG2370, whose sequence is provided in SEQ ID NO:4, comprises the vector backbone from the binary vector pBIN19 (GI1256363), T-DNA left and right border fragments, and, between border fragments, the CsVMV promoter sequence and a Nos termination sequence for controlling expression of the inserted gene, and the neomycin phosphotransferase (NPTII) gene, which confers kanamycin resistance, whose expression is controlled by the RE4 promoter (US Patent No. 6054635) and the G7 termination sequence. The *ANT1* fragment was cloned into SmaI/SpeI sites of pAG2370, inserted between the CsVMV promoter region, proximal to the 5' end of genomic fragment, and the Nos termination sequence, proximal to the 3' end of the genomic fragment. The pAG2370-*ANT1* construct was transformed into *Agrobacterium tumefaciens* by electroporation.

The pAG2370-*ANT1* construct described above was introduced into wild-type Micro-Tom plants via *Agrobacterium*-mediated transformation, essentially as described in Example 1. Briefly, explants were dissected from Micro-Tom seedlings. Explants were inoculated by soaking in the *Agrobacterium* suspension for 15 to 120 minutes, blotted on sterile filter paper to remove excess bacteria, and plated. Explants were co-cultivated in non-selective media for 2-4 days at 24°C with a 16-hour photoperiod, after which they were transferred to selective media (with kanamycin) and returned to the growth room. Explants were transferred to fresh medium every two weeks until shoots were 0.5 to 1 cm tall. Shoots were excised from the explants, placed on selective medium with kanamycin in Phytatrays (Sigma), and returned to the growth room for two to four weeks. Shoots were observed for rooting, and rooted shoots were out-planted to soil and acclimated to the greenhouse. The transformation process generated 64 independent T<sub>0</sub> events.

Morphological observations demonstrated that 45 transgenic plants displayed the *ANT1* purple color phenotype and were either partially or entirely purple. Tissue was collected from six T<sub>1</sub> plants showing the *ANT1* phenotype, and RT-PCR was carried out using wild type as a control. While no *ANT1* gene expression could be detected in the wild-type control, five out of the six plants displaying the *ANT1* phenotype over-expressed the *ANT1* transcript. The internal control experiments, using a constitutively expressed actin gene, showed that all samples had similar levels of the actin expression.

#### EXAMPLE 4

##### Confirmation of Phenotype/Genotype Association in *Arabidopsis*

In order to further confirm the association between the *ANT1* phenotype and the *ANT1* gene in plants other than Micro-Tom, the *ANT1* gene was introduced into and over-expressed in wild type *Arabidopsis thaliana*.

The pAG2370-*ANT1* construct described above was introduced into wild-type *Arabidopsis* plants via *Agrobacterium*-mediated transformation using standard vacuum infiltration methods. All infiltrated seeds were plated in selective media containing kanamycin, and kanamycin-resistant T<sub>1</sub> plants were transplanted to 72-cell flats. The transformation process generated 10 independent T<sub>0</sub> events, of which seven displayed the *ANT1* purple coloration phenotype in at least part of the plant. Tissue was collected from four T<sub>1</sub> plants showing the *ANT1* phenotype, and RT-PCR was carried out using wild type as a control. While no *ANT1* gene expression could be detected in the wild-type control, all plants displaying the *ANT1* phenotype over-expressed the *ANT1* transcript. The internal control experiments, using a constitutively expressed actin gene, showed that all samples had similar levels of the actin expression.

#### EXAMPLE 5

##### Confirmation of Phenotype/Genotype Association in Tobacco

In order to further confirm the association between the *ANT1* phenotype and the *ANT1* gene in plants other than Micro-Tom, the *ANT1* gene was introduced into and over-expressed in wild type *Nicotiana tabacum* (tobacco, Wisconsin-38 type).

The pAG2370-*ANT1* construct described above was introduced into wild-type tobacco plants via *Agrobacterium*-mediated transformation using essentially the following methods. In order to generate tobacco plants for transformation, tobacco seeds were germinated as follows: seeds were shaken about ten minutes on a lab shaker, in a solution

containing approximately 1.3% to 2.1% sodium hypochlorite and one drop of Tween-20 (Polyoxyethylenesorbitan monolaurate) per 100 milliliters. Seeds were then washed in sterile water and sterilely transferred to the surface of TbSG medium (4.3 g/l Murashige and Skoog salts, Phytotech; 1 ml/l MS vitamins, Sigma; 30 g/l sucrose; 8 g/l agar, Sigma; pH adjusted to ~5.8) in petri dishes or Phytatrays (Sigma), 10-50 seeds per vessel, and incubated in light at 25°C. Tobacco plants were dissected on sterile filter paper moistened with sterile, deionized water or liquid TbCo medium (4.3 g/l Murashige and Skoog salts, Phytotech; 1 ml/l MS vitamins, Sigma; 30 g/l sucrose; 200 mg/l  $\text{KH}_2\text{PO}_4$ ; 2 mg/l Indole-3-acetic acid; 0.25 mg/l Kinetin; 0 to 100  $\mu\text{M}$  Acetosyringone; 7 g/l Agar, Sigma; pH adjusted to 5.4-5.6). Explants with cut edges on all sides could be generated by cutting the leaf from the plant, dissecting out and discarding the midvein, and cutting the leaf lamina into 3 to 5 mm squares. Alternatively, discs could be cut from the lamina using a sterilized cork borer.

Explants were inoculated by soaking for 15-120 minutes in *Agrobacterium* suspension (OD<sub>600</sub> between 0.175 and 0.225) prepared with the pAG2370-*ANTI* construct, then blotted and plated on TbCo medium. Explants were co-cultivated 2-4 days at 24°C with a 16-hour photoperiod, and then transferred to Tb selective medium (4.3 g/l Murashige and Skoog salts; 1 ml/l Nitsch and Nitsch vitamins, Duchefa; 30 g/l sucrose; 0.5 to 2 mg/l 6-Benzylaminopurine; 0 to 1 mg/l Naphthylacetic Acid; 0 to 750 mg/l Carbenicillin; 0 to 300 mg/l Timentin; 0 to 500 mg/l Kanamycin; 7 to 8 g/l Agar, Sigma; pH adjusted to ~5.8) containing kanamycin and re-transferred every two weeks until shoots were 0.5 to 1 cm tall. Shoots were excised from the explants, placed on TbR medium (4.3 g/l Murashige and Skoog salts; 1 ml/l Nitsch and Nitsch vitamins, Duchefa; 30 g/l sucrose; 0 to 1 mg/l Indole-3-butyric acid; 0 to 1 mg/l Naphthylacetic Acid; 0 to 100 mg/l Carbenicillin; 0 to 200 mg/l Timentin; 0 to 100 mg/l Kanamycin; 7 to 8 g/l Agar, Sigma; pH adjusted to ~5.8.) with kanamycin in Phytatrays, and grown two to four weeks, after which time the rooted shoots were planted to soil.

The transformation process generated 89 independent T<sub>0</sub> events, of which 54 displayed the *ANTI* purple coloration phenotype in at least part of the plant. Tissue was collected from five T<sub>1</sub> plants showing the *ANTI* phenotype, and RT-PCR was carried out using wild type as a control. While no *ANTI* gene expression could be detected in the wild-type control, all plants displaying the *ANTI* phenotype over-expressed the *ANTI* transcript. The internal control experiments, using a constitutively expressed actin gene, showed that all samples had similar levels of the actin expression.

EXAMPLE 6Use of the *ANT1* gene as a transformation marker in tomato and tobacco

Having successfully recapitulated the *ANT1* phenotype in tomato and tobacco, as described above, we tested the utility of the *ANT1* gene for utility as a transformation marker, based on its characteristic purple color, in these species. We transformed tobacco and Micro-Tom explants with the pAG2370-*ANT1* vector, using methods described in the above Examples, grew the explants in the presence and absence of antibiotic (kanamycin), and compared transformation frequency based on rooting in the presence of antibiotic in the media to transformation frequency based on purple color. Results are shown in the Table below.

Table 1:

Transformation frequency of tobacco and tomato, based on antibiotic selection or color

Species	Kanamycin in media	# explants	Transformation frequency based on rooting in presence of antibiotic	Transformation frequency based on purple color
<u>Tobacco</u> Wiscons in	+	82	126 % *	80%
	-	60	-	45%
<u>Tomato</u> Micro-Tom	+	103	77%	54%
	-	52	-	6%

\* This number reflects multiple transgenic events per original explant. When callus initiation occurs at two or three distinct points on the original explant, each is dissected and tested for shoot regeneration.

The results indicated that the *ANT1* gene could be successfully used for screening of positive transformants in cultures of tomato and tobacco, and may be useful in other plants as well.

EXAMPLE 7Anthocyanin Analysis in Tomato and Tobacco Over-expressing *ANT1*

We performed liquid chromatography/mass spectrometry (LC/MS) analysis to characterize the anthocyanins produced in transgenic tomato and tobacco that mis-express *ANT1*.



Fresh tissue samples included leaves from wild-type micro-tom plants, from the original ACTTAG mutant line H000001624 ("*ANT1* micro-tom"), and from plants in which pAG2370-*ANT1* had been introduced ("recapitulated" *ANT1* micro-tom), as well as tobacco leaves from plants in which pAG2370-*ANT1* had been introduced ("*ANT1* tobacco"). Frozen tissue samples included leaves from *ANT1* micro-tom and from recapitulated *ANT1* micro-tom, and a pool of *in vitro* shoots of *ANT1* tomato.

To obtain tissue extracts, fresh or frozen plant material (0.5 g) was soaked for 2 hrs in 1ml of either 1% HCl/MeOH or 5% HOAc(aq). The extracted plant material was separated from the liquid phase by filtration or centrifugation and the extract was transferred to 2 ml vials for HPLC analysis.

The crude plant extract (10  $\mu$ l) was injected onto a Waters 2795 HPLC equipped with a Waters C-18, 3.5  $\mu$ m SymmetryShield column (4.6 x 150 mm). The extracts were eluted with a 30 minute mobile phase gradient of 5-35% ACN in 1.5% H<sub>3</sub>PO<sub>4</sub>(aq) with a flow rate of 1 ml/min. Compounds were detected at 520 nm using a Waters 996 photodiode array detector.

For LC/MS analysis, extracts (5  $\mu$ l) were chromatographed on a C-18 Symmetry column with a 0.1% formic acid(aq)/ACN mobile phase running at 0.3 ml/min. A Micromass Quattro Micro triple quadrupole mass spectrometer with an ES+ source was used to detect and analyze all components of the extract.

We compared the anthocyanin composition in the leaves of several *ANT1*, recapitulated *ANT1*, and wild-type micro-tom. All of the *ANT1* micro-tom plants were found to contain a similar mixture of at least nine different anthocyanins, although the ratios of the components varied. These anthocyanins appear to be elevated more than 100-fold compared to the wild-type (where they are almost undetectable), although concentrations differed from plant to plant, with the highest levels occurring in recapitulated lines. Mass spectral fragmentation of the nine anthocyanin molecular ions yielded one of three daughter ions at 303, 317, or 331 atomic mass units (AMU), indicating the presence of delphinidin, petunidin, and malvidin-type anthocyanidins (aglycones), respectively. With knowledge of the core structures, we realized that each anthocyanidin must be functionalized in the same three ways. Comparison of the molecular weights and MS fragmentation patterns of the tomato anthocyanins with common anthocyanin glycosylation motifs indicated that the *ANT1* tomato produces the anthocyanins listed in Table 2 below, and depicted in Figure 1. Anthocyanins designated (\*) in Table 2 have been reported in light stressed tomatoes (Bory et al, The Plant Cell

(2002) 14:2509-2526). Presence of the remaining six molecules in tomato has not been reported previously.

Table 2

Substitution Pattern (Fig. 1)	Anthocyanin	Molecular Ion (M <sup>+</sup> )
R <sub>1</sub> =OH, R <sub>2</sub> =OH, R <sub>3</sub> =H	Delphinidin 3-rutinoside-5-glucoside	773.4
R <sub>1</sub> =OH, R <sub>2</sub> =OH, R <sub>3</sub> =Coumarate	Delphinidin 3-(coumaroyl)rutinoside-5-glucoside	919.6
R <sub>1</sub> =OH, R <sub>2</sub> =OH, R <sub>3</sub> =Caffeate	Delphinidin 3-(caffeoyl)rutinoside-5-glucoside	935.6
R <sub>1</sub> =OMe, R <sub>2</sub> =OH, R <sub>3</sub> =H	Petunidin 3-rutinoside-5-glucoside	787.4
R <sub>1</sub> =OMe, R <sub>2</sub> =OH, R <sub>3</sub> =Coumarate	Petunidin 3-(coumaroyl)rutinoside-5-glucoside (*)	933.6
R <sub>1</sub> =OMe, R <sub>2</sub> =OH, R <sub>3</sub> =Caffeate	Petunidin 3-(caffeoyl)rutinoside-5-glucoside (*)	949.6
R <sub>1</sub> =OMe, R <sub>2</sub> =OMe, R <sub>3</sub> =H	Malvidin 3-rutinoside-5-glucoside	801.4
R <sub>1</sub> =OMe, R <sub>2</sub> =OMe, R <sub>3</sub> =Coumarate	Malvidin 3-(coumaroyl)rutinoside-5-glucoside (*)	947.6
R <sub>1</sub> =OMe, R <sub>2</sub> =OMe, R <sub>3</sub> =Caffeate	Malvidin 3-(caffeoyl)rutinoside-5-glucoside	963.6

LC/MS analysis of *ANT1* tobacco indicated two major anthocyanins with molecular weights of 449 and 595. Both components fragmented to give daughter ions at 287 amu, indicating a cyanidin type nucleus. Since cyanidins commonly occur as glycosides, we deduced that *ANT1* tobacco contains mainly cyanidin-3-glucoside and cyanidin-3-rutinoside (Figure 2).

### EXAMPLE 8

#### Isoflavone Analysis in Tomato Over-expressing *ANT1*

Because isoflavones are derived from the phenylpropanoid pathway that also gives rise to anthocyanins, we analyzed fruits and leaves from an *ANT1* and wild type micro-tomato for daidzein, genistein, and glycitein.

Isoflavone analysis was performed by Covance Laboratories Inc (Madison WI) according to the method of Seo and Morr (1984, J Agric Food Chem 32: 530-533) and Petterson and Kiessling (1984, J Assoc Off Anal Chem, 67:503-506). The detection limit of the analysis was 1.0 mg/100g for glycitein, daidzein and genistein. As expected, wild type tomato had no detectable isoflavones in either leaves or fruit. However, leaves of

*ANT1* micro-tomato produced detectable levels of glycitein at nearly twice the detection limit. Leaves of *ANT1* produced glycitein at 1.91 mg/100g compared to <1.00 mg/100g in the wild type.